

Cat. No.: <u>ETAQ-101</u>

#### E I ERNAL LIFE SCIENCE

# Taq DNA Polymerase Mastermix (2X)

(ETERBIO-EZ-ETAQ-101)

Lot:
Expiry Date: <u>yy</u> / <u>mm</u> / <u>dd</u>
g. 200g
Store at -20°C

#### DESCRIPTION

Taq DNA Polymerase Mastermix (2X) is a ready-to-use 2X solution containing Taq DNA polymerase, dNTPs, MgCl<sub>2</sub> and reaction buffers at optimal concentrations for PCR. Simply add primers, template, and water to amplify the target sequence and other molecular biology applications. This formulation not only saves valuable time, but also reduces number of pipetting and reagent handling errors.

## Composition of the Taq DNA Polymerase Mastermix (2X)

 $0.2 \text{ U/}\mu l$  Taq DNA polymerase, reaction buffer, 3 mM MgCl<sub>2</sub>, 0.4 mM of each dNTP (dATP, dCTP, dGTP and dTTP).


# **USAGE**

- 1. Gently vortex and briefly centrifuge Taq DNA Polymerase Mastermix (2X) after thawing.
- 2. Set up each reaction as follows:

Component	50 μl reaction	Final Concentration
Mastermix (2X)	25 μ1	1X
Primer A	Variable	0.1–1.0 μM
Primer B	Variable	0.1–1.0 μM
Template DNA	Variable	< 1.0 μg
Nuclease-free water	to 50 μ1	

- 3. Gently mix the solution a few times and spin down.
- Perform PCR using the recommended thermal cycling conditions outlined below:
  (For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair)

Step	Temperature	Time
Initial Denaturation	95°C	30 seconds
30 Cycles	95°C	15-30 seconds
	45-68°C	15-60 seconds
	72°C	1 minute/kb
Final Extension	72°C	5 minutes
Hold	4-10°C	





## **Endonuclease Assay**

No conversion of covalently closed circular DNA to nicked form was detected after incubation of 1X *Taq* DNA Polymerase Mastermix with 1 µg of supercoiled plasmid DNA (pUC19) in for 4 hours at 37 °C.

## **Exonuclease Assay**

No degradation of DNA was observed after incubation of 1 µg of lambda DNA/*Hin*dIII fragments in 1X *Taq* DNA Polymerase Mastermix for 4 hours at 37°C.

## **Functional Assay**

Good performance in PCR was tested for amplification of 1.8 kb gene.

# For Research Use Only